

# Some directions in Perfect Simulation

Laird Breyer,  
<mailto:laird@lbreyer.com>

joint work with **Gareth Roberts** (Lancaster)

The work presented here has involved the following institutions:



University of Lancaster



Aalborg University



Università di Roma Tre



# Aims of this talk

In this talk, I shall present loosely some recent ideas related to Perfect Simulation, including:

# Aims of this talk

In this talk, I shall present loosely some recent ideas related to Perfect Simulation, including:

- what Read-Once CFTP means

# Aims of this talk

In this talk, I shall present loosely some recent ideas related to Perfect Simulation, including:

- what Read-Once CFTP means
- some old and new coupling constructions

# Aims of this talk

In this talk, I shall present loosely some recent ideas related to Perfect Simulation, including:

- what Read-Once CFTP means
- some old and new coupling constructions
- an example of perfect simulation

# Aims of this talk

In this talk, I shall present loosely some recent ideas related to Perfect Simulation, including:

- what Read-Once CFTP means
- some old and new coupling constructions
- an example of perfect simulation

PS is still at the beginning, and new methods are discovered regularly...


# MCMC and Perfect Samples

In MCMC, we commonly have two distinct goals:

# MCMC and Perfect Samples

In MCMC, we commonly have two distinct goals:

Simulation



Integration


$$\mathbb{P}(A \mid B) = \int_A \mathbb{P}(B \mid \omega) \mathbb{P}(d\omega) / \mathbb{P}(B)$$



# MCMC and Perfect Samples

In MCMC, we commonly have two distinct goals:

Simulation



Integration


$$\mathbb{P}(A \mid B) = \int_A \mathbb{P}(B \mid \omega) \mathbb{P}(d\omega) / \mathbb{P}(B)$$

PS solves the simulation problem completely,

# MCMC and Perfect Samples

In MCMC, we commonly have two distinct goals:

Simulation



Integration

$$\mathbb{P}(A \mid B) = \int_A \mathbb{P}(B \mid \omega) \mathbb{P}(d\omega) / \mathbb{P}(B)$$

PS solves the simulation problem completely, and reduces integration problems to Classical Statistics.

# Building Markov chains

There are several equivalent definitions for Markov chains. For PS, we use a computational definition:

A sequence  $X_1, X_2, \dots$  is a Markov chain if there exists an IID series of random functions  $F_1(x), F_2(x), \dots$  such that


$$F_1(X_1) = X_2, \quad F_2(X_2) = X_3, \quad \dots, \quad F_k(X_k) = X_{k+1}$$

# Building Markov chains

There are several equivalent definitions for Markov chains. For PS, we use a computational definition:

A sequence  $X_1, X_2, \dots$  is a Markov chain if there exists an IID series of random functions  $F_1(x), F_2(x), \dots$  such that

$$F_1(X_1) = X_2, \quad F_2(X_2) = X_3, \quad \dots, \quad F_k(X_k) = X_{k+1}$$



# The stationary density

Consider the chain over a larger time scale: the stationary density  $\pi$  satisfies  $\pi P = \pi$ , where


$$P(x, dy) = \mathbb{P}(F_t(x) \in dy) \quad \text{for all } t.$$

# The stationary density

Consider the chain over a larger time scale: the stationary density  $\pi$  satisfies  $\pi P = \pi$ , where

$$P(x, dy) = \mathbb{P}(F_t(x) \in dy) \quad \text{for all } t.$$

Here is a path:




# The stationary density

Consider the chain over a larger time scale: the stationary density  $\pi$  satisfies  $\pi P = \pi$ , where

$$P(x, dy) = \mathbb{P}(F_t(x) \in dy) \quad \text{for all } t.$$

After reordering the occupied states, we get:




# Read Once Perfect Simulation

Wilson (1999) showed how to recognize random times  $T_2, T_3, \dots$  when the chain is *exactly* in equilibrium:




# Read Once Perfect Simulation

Wilson (1999) showed how to recognize random times  $T_2, T_3, \dots$  when the chain is *exactly* in equilibrium:



# Read Once Perfect Simulation

Wilson (1999) showed how to recognize random times  $T_2, T_3, \dots$  when the chain is *exactly* in equilibrium:



# How to find the perfect samples

- Trace all possible paths under the mappings  $F_1, F_2, \dots$ , looking for *coalescence* within  $m$  steps:
- Let  $G_1(x), G_2(x), \dots$  be defined for all  $x$  by:

$$G_1(x) = F_m(F_{m-1}(\cdots F_1(x) \cdots))$$


$$G_2(x) = F_{2m}(F_{2m-1}(\cdots F_{m+1}(x) \cdots))$$

and set

$$T_1 = \min\{u \geq 0 : G_u(x) = G_u(y) \text{ for all } x, y\}$$

$$T_{k+1} = \min\{u \geq T_k : G_u(x) = G_u(y) \text{ for all } x, y\}$$

# How to find the perfect samples



# The fundamental identity

Let  $\epsilon = \mathbb{P}(T_1 \leq 1)$  and write

$$\begin{aligned}\mathbb{P}(G_1(x) \in dy) &= (1 - \epsilon)\mathbb{P}(G_1(x) \in dy \mid T_1 > 1) + \epsilon\mathbb{P}(G_1(x) \in dy \mid T_1 \leq 1) \\ &= (1 - \epsilon)Q(x, dy) + \epsilon\mu(dy).\end{aligned}$$

Then  $\pi = \epsilon \sum_{s=0}^{\infty} (1 - \epsilon)^s \mu Q^s$ .

# The fundamental identity

Let  $\epsilon = \mathbb{P}(T_1 \leq 1)$  and write

$$\begin{aligned}\mathbb{P}(G_1(x) \in dy) &= (1 - \epsilon)\mathbb{P}(G_1(x) \in dy \mid T_1 > 1) + \epsilon\mathbb{P}(G_1(x) \in dy \mid T_1 \leq 1) \\ &= (1 - \epsilon)Q(x, dy) + \epsilon\mu(dy).\end{aligned}$$

Then  $\pi = \epsilon \sum_{s=0}^{\infty} (1 - \epsilon)^s \mu Q^s$ .

*Proof.* Using stationarity  $\pi P = \pi$ , we have

$$(1 - \epsilon)^k \pi Q^k = (1 - \epsilon)^{k-1} \pi (P - \epsilon\mu) Q^{k-1}$$

# The fundamental identity

Let  $\epsilon = \mathbb{P}(T_1 \leq 1)$  and write

$$\begin{aligned}\mathbb{P}(G_1(x) \in dy) &= (1 - \epsilon)\mathbb{P}(G_1(x) \in dy \mid T_1 > 1) + \epsilon\mathbb{P}(G_1(x) \in dy \mid T_1 \leq 1) \\ &= (1 - \epsilon)Q(x, dy) + \epsilon\mu(dy).\end{aligned}$$

Then  $\pi = \epsilon \sum_{s=0}^{\infty} (1 - \epsilon)^s \mu Q^s$ .

*Proof.* Using stationarity  $\pi P = \pi$ , we have

$$\begin{aligned}(1 - \epsilon)^k \pi Q^k &= (1 - \epsilon)^{k-1} \pi (P - \epsilon \mu) Q^{k-1} \\ &= (1 - \epsilon)^{k-1} \pi Q^{k-1} - \epsilon (1 - \epsilon)^{k-1} \mu Q^{k-1}\end{aligned}$$

# The fundamental identity

Let  $\epsilon = \mathbb{P}(T_1 \leq 1)$  and write

$$\begin{aligned}\mathbb{P}(G_1(x) \in dy) &= (1 - \epsilon)\mathbb{P}(G_1(x) \in dy \mid T_1 > 1) + \epsilon\mathbb{P}(G_1(x) \in dy \mid T_1 \leq 1) \\ &= (1 - \epsilon)Q(x, dy) + \epsilon\mu(dy).\end{aligned}$$

Then  $\pi = \epsilon \sum_{s=0}^{\infty} (1 - \epsilon)^s \mu Q^s$ .

*Proof.* Using stationarity  $\pi P = \pi$ , we have

$$\begin{aligned}(1 - \epsilon)^k \pi Q^k &= (1 - \epsilon)^{k-1} \pi (P - \epsilon \mu) Q^{k-1} \\ &= (1 - \epsilon)^{k-1} \pi Q^{k-1} - \epsilon (1 - \epsilon)^{k-1} \mu Q^{k-1} \\ &= \dots = \\ &= \pi - \epsilon \sum_{s=1}^k (1 - \epsilon)^{k-s} \mu Q^{k-s}.\end{aligned}$$



# The fundamental identity

Let  $\epsilon = \mathbb{P}(T_1 \leq 1)$  and write

$$\begin{aligned}\mathbb{P}(G_1(x) \in dy) &= (1 - \epsilon)\mathbb{P}(G_1(x) \in dy \mid T_1 > 1) + \epsilon\mathbb{P}(G_1(x) \in dy \mid T_1 \leq 1) \\ &= (1 - \epsilon)Q(x, dy) + \epsilon\mu(dy).\end{aligned}$$

Then  $\pi = \epsilon \sum_{s=0}^{\infty} (1 - \epsilon)^s \mu Q^s$ .

*Proof.* Using stationarity  $\pi P = \pi$ , we have

$$\begin{aligned}(1 - \epsilon)^k \pi Q^k &= (1 - \epsilon)^{k-1} \pi (P - \epsilon \mu) Q^{k-1} \\ &= (1 - \epsilon)^{k-1} \pi Q^{k-1} - \epsilon (1 - \epsilon)^{k-1} \mu Q^{k-1} \\ &= \dots = \\ &= \pi - \epsilon \sum_{s=1}^k (1 - \epsilon)^{k-s} \mu Q^{k-s}.\end{aligned}$$

Let  $k \rightarrow \infty$ , then left side goes to zero.

# How to ensure coupling occurs

Again assume that we have an IID sequence  $F_1, F_2, \dots$  of random maps satisfying

$$P(x, dy) = \mathbb{P}(F_t(x) \in dy) \quad \text{for all } t. \quad (1)$$

With  $F_t(x)$  arbitrary, coalescence of Markov chain sample paths need not ever happen. We must modify the maps subject to preserving the condition (??).

# Modifying the updates

- Normally at each iteration, we use a randomly generated function  $F_t(x)$  such that  $X_{t+1} = F_t(X_t)$ .

# Modifying the updates

- Normally at each iteration, we use a randomly generated function  $F_t(x)$  such that  $X_{t+1} = F_t(X_t)$ .
- If  $F_t(x) = F_t(x')$  holds for some  $x, x'$ , then there is the *possibility* of coupling.

# Modifying the updates


- Normally at each iteration, we use a randomly generated function  $F_t(x)$  such that  $X_{t+1} = F_t(X_t)$ .
- If  $F_t(x) = F_t(x')$  holds for some  $x, x'$ , then there is the *possibility* of coupling.
- **Strategy:** From  $F$ , generate a new random function  $\mathcal{C}_Y(F)$  which has a higher chance of coupling.

# Modifying the updates


- Normally at each iteration, we use a randomly generated function  $F_t(x)$  such that  $X_{t+1} = F_t(X_t)$ .
- If  $F_t(x) = F_t(x')$  holds for some  $x, x'$ , then there is the *possibility* of coupling.
- **Strategy:** From  $F$ , generate a new random function  $\mathcal{C}_Y(F)$  which has a higher chance of coupling.
- **Definition:** Let  $Y$  be independent of  $F$ , with  $\mathbb{P}(Y = y) = q(y)$  say.

$$\mathcal{C}_Y(F)(x) = \begin{cases} Y & \text{if } \frac{p(x,Y)q(F(x))}{p(x,F(x))q(Y)} > U[0, 1] \\ F(x) & \text{otherwise.} \end{cases}$$


# Modifying the updates




$f$



$G_Y(f)$




$G_Y(G_Y(f))$




$G_Y(G_Y(G_Y(G_Y(G_Y(f)))))$


# Modifying the updates




$f$



$G_Y(f)$



$G_Y(G_Y(f))$



$G_Y(G_Y(G_Y(G_Y(G_Y(f)))))$

We call the triangles catalysts. Click [here](#).



# Why it works

- In the formula

$$\mathcal{C}_Y(F)(x) = \begin{cases} Y & \text{if } \frac{p(x,Y)q(F(x))}{p(x,F(x))q(Y)} > U[0, 1] \\ F(x) & \text{otherwise.} \end{cases}$$

fix  $x$  and define  $\Pi(y) = p(x, y)$ ,  $X = F(x)$  and  $X' = \mathcal{C}_Y(F)(x)$ .

# Why it works

- In the formula

$$\mathcal{C}_Y(F)(x) = \begin{cases} Y & \text{if } \frac{p(x,Y)q(F(x))}{p(x,F(x))q(Y)} > U[0, 1] \\ F(x) & \text{otherwise.} \end{cases}$$

fix  $x$  and define  $\Pi(y) = p(x, y)$ ,  $X = F(x)$  and  $X' = \mathcal{C}_Y(F)(x)$ .

- We get

$$X' = \begin{cases} Y & \text{if } \frac{\Pi(Y)q(X)}{\Pi(X)q(Y)} > U[0, 1] \\ X & \text{otherwise} \end{cases}$$

# Why it works

- In the formula

$$\mathcal{C}_Y(F)(x) = \begin{cases} Y & \text{if } \frac{p(x,Y)q(F(x))}{p(x,F(x))q(Y)} > U[0, 1] \\ F(x) & \text{otherwise.} \end{cases}$$


fix  $x$  and define  $\Pi(y) = p(x, y)$ ,  $X = F(x)$  and  $X' = \mathcal{C}_Y(F)(x)$ .

- We get

$$X' = \begin{cases} Y & \text{if } \frac{\Pi(Y)q(X)}{\Pi(X)q(Y)} > U[0, 1] \\ X & \text{otherwise} \end{cases}$$

- Therefore, since  $\Pi(dy) = \mathbb{P}(X \in dy)$  by assumption, it follows that  $\mathbb{P}(X' \in dy) = \Pi(dy)$  also.

# Extensions



- At every update, let the catalyst  $Y_t$  come from an independent Markov chain with identical transition probabilities. Great for Gibbs samplers!
- We have the formula

$$\|\mathbb{P}(X_t \in \cdot) - \pi\|_{\text{TV}} \leq \mathbb{E} \prod_{s=0}^t \left( 1 - \frac{p(X_s, Y_{s+1})p(Y_s, X_{s+1})}{p(X_s, X_{s+1})p(Y_s, Y_{s+1})} \right)_+,$$

# Pump Example (Autogamma)

- Let  $x = (\beta, \lambda_1, \dots, \lambda_{10})$ , simulate

$$\pi(x) = \exp \left\{ (10\alpha + \gamma - 1) \log \beta - \delta \beta \right. \\ \left. + \sum_{k=1}^{10} \left( (s_k + \alpha - 1) \log \lambda_k - (\beta + t_k) \lambda_k \right) \right\},$$

# Pump Example (Autogamma)

- Let  $x = (\beta, \lambda_1, \dots, \lambda_{10})$ , simulate

$$\pi(x) = \exp \left\{ (10\alpha + \gamma - 1) \log \beta - \delta \beta \right. \\ \left. + \sum_{k=1}^{10} \left( (s_k + \alpha - 1) \log \lambda_k - (\beta + t_k) \lambda_k \right) \right\},$$

- **Gibbs sampler:** One sweep is  $f : (\beta, \lambda_1, \dots, \lambda_{10}) \mapsto (\beta', \lambda'_1, \dots, \lambda'_{10})$ , where

$$\beta' \sim \pi_0(\cdot \mid \lambda_1, \dots, \lambda_{10}) = \Gamma(\gamma + 10\alpha, \delta + \sum_{k=1}^{10} \lambda_k),$$

$$\lambda'_k \sim \pi_k(\cdot \mid \beta') = \Gamma(\alpha + s_k, \beta' + t_k), \quad k = 1, \dots, 10.$$

# Pump Example: coupling

- Transition density is

$$p(\beta, \lambda_1, \dots, \lambda_{10}; b, l_1, \dots, l_{10}) = \pi_0(b \mid \lambda_1, \dots, \lambda_{10}) \prod_{k=1}^{10} \pi_k(l_k \mid b).$$

# Pump Example: coupling

- Transition density is

$$p(\beta, \lambda_1, \dots, \lambda_{10}; b, l_1, \dots, l_{10}) = \pi_0(b \mid \lambda_1, \dots, \lambda_{10}) \prod_{k=1}^{10} \pi_k(l_k \mid b).$$

- Take  $Y \sim q$  where

$$q(y_0, y_1, \dots, y_{10}) = \pi_0(y_0 \mid \lambda_1^*, \dots, \lambda_{10}^*) \prod_{k=1}^{10} \pi_k(y_k \mid y_0),$$



# Pump Example: coupling

- After simplification,

$$\mathcal{C}_Y(f)(x) = \begin{cases} Y & \text{if } \exp\left((\beta' - Y_0)(|\lambda| - |\lambda^*|)\right) > \xi, \\ (\beta', \lambda'_1, \dots, \lambda'_{10}) & \text{otherwise.} \end{cases}$$

# Pump Example: coupling

- After simplification,

$$\mathcal{C}_Y(f)(x) = \begin{cases} Y & \text{if } \exp\left((\beta' - Y_0)(|\lambda| - |\lambda^*|)\right) > \xi, \\ (\beta', \lambda'_1, \dots, \lambda'_{10}) & \text{otherwise.} \end{cases}$$

- Thus

$$\text{Basin}(Y, f, \xi) = \left\{ x : \left( \psi_0 - (\delta + |\lambda|)Y_0 \right) \left( |\lambda| - |\lambda^*| \right) > \log \xi \right\},$$

# Pump Example: coupling

- After simplification,

$$\text{Basin}(Y, f, \xi) = \left\{ x : |\lambda| \in \left[ \frac{-b - \sqrt{b^2 - 4ac}}{2a}, \frac{-b + \sqrt{b^2 - 4ac}}{2a} \right] \right\},$$

where  $a = Y_0$ ,  $b = \log \xi - Y_0(|\lambda^*| - \delta)$  and  $c = \delta(\log \xi - Y_0|\lambda^*|) - \psi_0$ .

# Conclusions

- If we can recognize the basins, we can also recognize when the maps  $G_u$  coalesce.

# Conclusions

- If we can recognize the basins, we can also recognize when the maps  $G_u$  coalesce.
- If we can recognize map coalescence, we can also do Perfect Simulation.

# Conclusions

- If we can recognize the basins, we can also recognize when the maps  $G_u$  coalesce.
- If we can recognize map coalescence, we can also do Perfect Simulation.
- You can find a simulation by clicking [here](#).

# Learning more

Preprints and tutorials may be found at the following two sites:

- <http://www.dimacs.rutgers.edu/~dbwilson/exact.html>
- <http://www.lbreyer.com>